

Math 33A – Week 1

Written by Victoria Kala

April 2, 2019

Name: KEY

1. Consider the system

$$\begin{aligned} 2x - y &= 7 \\ -x + \frac{1}{2}y &= 10 \end{aligned}$$

(a) Solve the system using elimination or substitution.

elimination: $2x - y = 7$
 (multiply 2nd equation by 2) $\frac{-2x + y = 20}{0 + 0 = 27}$
 $0 = 27$ **no solution**

(b) How many solutions does this system have? *none*

(c) Solve using Gaussian or Gauss-Jordan elimination and show you have the same result as part (a).

$$\left(\begin{array}{cc|c} 2 & -1 & 7 \\ -1 & \frac{1}{2} & 10 \end{array} \right) \xrightarrow{R_1 \leftrightarrow R_2} \left(\begin{array}{cc|c} -1 & \frac{1}{2} & 10 \\ 2 & -1 & 7 \end{array} \right) \xrightarrow{2R_1 + R_2 \rightarrow R_2} \left(\begin{array}{cc|c} -1 & \frac{1}{2} & 10 \\ 0 & 0 & 27 \end{array} \right) \leftarrow \text{this last row says } 0 = 27 \Rightarrow \text{no solution, same as (a)}$$

2. Consider the system

$$\begin{aligned} x + 2y &= 13 \\ 4x + 8y &= 52 \end{aligned}$$

(a) Solve the system using elimination or substitution.

elimination: $-4x - 8y = -52$ let $y = t \rightarrow x + 2t = 13$
 (multiply 1st equation by -4) $\frac{4x + 8y = 52}{0 + 0 = 0}$
 $0 = 0$ infinitely many solutions
 $\boxed{\begin{matrix} x = 13 - 2t \\ y = t \end{matrix}}$ where $t \in \mathbb{R}$

(b) How many solutions does this system have? *infinitely many*

(c) Solve using Gaussian or Gauss-Jordan elimination and show you have the same result as part (a).

$$\left(\begin{array}{cc|c} 1 & 2 & 13 \\ 4 & 8 & 52 \end{array} \right) \xrightarrow{-4R_1 + R_2 \rightarrow R_2} \left(\begin{array}{cc|c} 1 & 2 & 13 \\ 0 & 0 & 0 \end{array} \right) \leftarrow \text{this last row says } 0 = 0 \Rightarrow \text{infinitely many solutions}$$

First row: let $y = t$
 $x + 2y = 13 \rightarrow \boxed{\begin{matrix} x = 13 - 2t \text{ or } (13 - 2t) \\ y = t \\ \text{where } t \in \mathbb{R} \end{matrix}}$
 Same as (a)

3. Consider the system

$$\begin{aligned}x + 3y &= 21 \\ 5x - y &= -7\end{aligned}$$

(a) Solve the system using elimination or substitution.

elimination:

$$\begin{array}{r} X + 3y = 21 \\ 15X - 3y = -21 \\ \hline 16X = 0 \\ X = 0 \end{array} \quad \begin{array}{l} \curvearrowright \\ 0 + 3y = 21 \\ y = 7 \end{array} \quad \boxed{\begin{array}{l} X = 0 \\ y = 7 \end{array}}$$

(multiply 2nd row by 3)

(b) How many solutions does this system have? *one*

(c) Solve using Gaussian or Gauss-Jordan elimination and show you have the same result as part (a).

$$\begin{pmatrix} 1 & 3 & | & 21 \\ 5 & -1 & | & -7 \end{pmatrix} \xrightarrow{-5R_1 + R_2 \rightarrow R_2} \begin{pmatrix} 1 & 3 & | & 21 \\ 0 & -16 & | & -112 \end{pmatrix} \xrightarrow{-\frac{1}{16}R_2 \rightarrow R_2} \begin{pmatrix} 1 & 3 & | & 21 \\ 0 & 1 & | & 7 \end{pmatrix} \xrightarrow{-3R_2 + R_1 \rightarrow R_1} \begin{pmatrix} 1 & 0 & | & 0 \\ 0 & 1 & | & 7 \end{pmatrix}$$

linear

$$\boxed{\begin{array}{l} X = 0 \\ y = 7 \end{array}} \text{ or } \begin{pmatrix} 0 \\ 7 \end{pmatrix} \text{ same as (a)}$$

4. (a) Is it possible for a system of ^{linear} equations to have exactly two solutions? *no*

(b) How many solutions can a linear system have? (*Hint*: Three possibilities)

0, 1, or ∞ many

(c) Using Gaussian or Gauss-Jordan elimination, how do you know when you have no solution?

0 = nonzero #

(d) Using Gaussian or Gauss-Jordan elimination, how do you know when you have infinitely many solutions?

* row of zeros (in square systems, not necessarily true in nonsquare systems)

* column with "missing pivot"

↑ pivot is leading entry in row