

Problem 1 Let \mathcal{D} be the region in \mathbb{R}^3 bounded by the surfaces (described in cylindrical coordinates) $\theta = z$, $\theta = z + \pi/2$, $z = 0$, and $z = \pi/2$.

- (a) Sketch the region \mathcal{D} .
- (b) Find the volume of \mathcal{D} .
- (c) Compute $\iiint_{\mathcal{D}} xyz \, dV$.

Problem 2 For each $h \in [0, 1]$, let \mathcal{R}_h be the region in \mathbb{R}^3 bounded by the surfaces $z = 0$, $z = h$, and $x^2 + y^2 + z^2 = 1$.

- (a) Sketch the regions \mathcal{R}_1 and $\mathcal{R}_{\frac{1}{2}}$.
- (b) Let $f(h) = \iiint_{\mathcal{R}_h} dV$. Explain why $f'(h) = \pi(1 - h^2)$.
- (c) Find $f(h)$ by evaluating the integral $\iiint_{\mathcal{R}_h} dV$ using cylindrical coordinates.
- (d) Find $f(h)$ by evaluating the integral $\iiint_{\mathcal{R}_h} dV$ using spherical coordinates.
- (e) Evaluate

$$\iiint_{\mathcal{R}_{1/2}} \frac{z}{\sqrt{x^2 + y^2 + z^2}} \, dV.$$

Problem 3 Let \mathcal{E} be the region defined by $x \geq 0$, $y \leq x$, $z \geq 0$, $x^2 + y^2 \leq 4$, and $z \leq y/x$. Evaluate

$$\iiint_{\mathcal{E}} (z - x^2 - y^2) \, dV.$$

Problem 4 Let \mathcal{E} be the region in the first octant bounded by $x^2 + y^2 + z^2 = 4$.

- (a) Evaluate

$$\iiint_{\mathcal{E}} (x - y + 3z) \, dV.$$

- (b) Evaluate

$$\iiint_{\mathcal{E}} (x^2 + y^2 - z^2) \, dV.$$

- (c) Griff makes a piece of Jell-O® whose shape is \mathcal{E} . Because Griff isn't very good at cooking, the density of her Jello-O® is very uneven: it is given by the function $\delta(x, y, z) = \frac{x^2 + y^2}{x^2 + y^2 + z^2}$. Find the mass of Griff's Jell-O®.

Problem 5 Let \mathcal{S} be the region bounded by $x = -1$, $x = 1$, $y + z = 0$, $y - z = 0$, and $y^2 + z^2 = 2$. Find the volume of \mathcal{S} (Hint: sketch what \mathcal{S} looks like and think about which coordinate system to use).